

Gabapentin Tablet 300 mg

Product Information :

Pharmaceutical form : Tablet

Category : Prescription Medicine

Pack : 1x15/Tabs

Brand name : Gabapentin Tablets

Content : Gabapentin (300mg)

Therapeutic Use : Treatment of partial seizures with and without secondary generalisation



Product Description :

What is Gabapentin Tablet 300 mg?

Gabapentin Tablet 300 mg is used in combination with other drugs to help people with epilepsy control some types of seizures. Gabapentin tablets can also be used to treat postherpetic neuralgia pain (the burning, stabbing pain or aches). Restless legs syndrome is treated with gabapentin extended-release tablet (a situation that causes discomfort in the legs). It belongs to the anticonvulsant class of medicines. Gabapentin cures seizures by reducing aberrant brain excitation. It can improve PHN discomfort by altering how the body perceives pain. The specific mechanism of action of gabapentin in the treatment of restless legs syndrome is unknown.

How to Use

- Gabapentin is available in the form of a capsule, tablet, extended-release tablet, and liquid. Gabapentin medications are normally taken 3 times a day with a full glass of water, both with and without meals.
- These drugs should be taken at regular intervals during the day and night, with no more

than 12 hours of intervals should pass between the doses.

- Do not cut, chew, or break the Gabapentin Tablet 300 mg; they must be swallowed whole.
- If your doctor has prescribed half of a standard tablet as part of your treatment, properly break the tablet along the mark. Use the remaining half-tablet in your next dose. Discard any half-tablets you haven't used within a few days of splitting them properly.
- If you're receiving gabapentin in the treatment of seizures or PHN, your doctor would most likely start you on a low dose and progressively raise it as necessary to treat your situation.
- Gabapentin can help you manage your illness, but it won't make it go away. Even though you feel better, keep taking gabapentin. You may have withdrawal problems such as anxiety, trouble falling asleep, nausea, discomfort, and sweating if you suddenly stop using gabapentin tablets.
- Gabapentin is also used in the treatment and prevention of hot flashes (sudden high feelings of heat and perspiration) in women who have been treated for breast cancer or have gone through menopause (the end of monthly menstrual cycles).

Benefits of Gabapentin Tablet

In Neuropathic Pain Treatment

Gabapentin Tablets are used in the treatment of long-term pain caused by nerve damage, such as that caused by diabetes, shingles, or a spinal cord injury. It relieves pain and its effects, such as mood swings, sleep issues, and exhaustion. It is considered to function by interacting with pain signals sent to the brain from injured nerves. Taking regular this medication will enhance your personal and social functioning, as well as your general quality of life. It takes some weeks to work, so take it on a regular basis even if you don't think it's helping. Even if your difficulties have subsided, you should continue to take the medication until your doctor recommends otherwise.

In Epilepsy/Seizures Treatment

Gabapentin Tablet 300 mg relates to the anticonvulsant drug class, which is used in the treatment of some types of epilepsy in adults. It is considered to function by slowing down the nerve impulses that produce seizures in the brain. Confusion, uncontrolled jerking

motions, lack of awareness, and anxiety or worry are some of the symptoms it helps with. It will give you greater confidence in your daily activities and may enable you to participate in things that you would instead be disallowed or afraid to perform. This treatment takes a few weeks to work, and you may still suffer seizures during that period. Even if you feel fine, do not stop taking this prescription until your doctor tells you to. Don't miss the dose.

Precaution - Gabapentin Tablet 300 mg

- If you have an allergy to gabapentin, any other drugs, or any of the inactive components in the gabapentin you plan to use, notify your doctor.
- You should be aware that gabapentin comes in a variety of forms that can be given for a variety of purposes. Check with your doctor to be sure you're not getting more than one gabapentin medicine.
- Take antacids for at least two hours before consuming gabapentin tablets, capsules, or solutions if you are using them.
- Talk to your doctor that you are receiving gabapentin if you are getting surgery, including dental surgery.
- Tell your doctor about any prescription and OTC medications, vitamins, nutraceuticals, and herbal products you're using or planning to use.
- If you have or have previously had lung or kidney problems, talk to your doctor.
- If you have a pregnancy, planning to become pregnant, or breastfeeding, notify your doctor. Call your doctor when you become pregnant while receiving Gabapentin Tablet 300 mg.
- You should be aware that this medicine may induce drowsiness or dizziness, as well as slowed thinking and incoordination. Do not drive or operate heavy machinery until you have a better understanding of how this drug affects you.
- Keep in mind that drinking alcohol can exacerbate the drowsiness brought on by this drug.
- While using gabapentin to treat epilepsy, mental disorder, or other diseases, you should be aware that your mental state may alter in unexpected ways, and you may feel suicidal (thinking about injuring or killing oneself, or planning or attempting to do so). If you, your family, or your caretaker notice any of the symptoms listed, contact your doctor

immediately.

Side Effects of Gabapentin

The majorities of side effects are minor and will go away as your body adapts to the medication. If they don't go away or you're concerned about them, see your doctor.

- It's possible that you'll experience drowsiness, dizziness, incoordination, weariness, blurred vision, strange eye movements, or tremors. Inform your doctor immediately if any of these side effects continue or worsen.
- Remember that your doctor provided Gabapentin Tablet 300 mg because he or she believes the advantage to you exceeds the risk of adverse effects. The majority of people who take this medicine do not have any substantial adverse effects.
- If any of these uncommon but significant side effects develop, contact your doctor immediately: swelling of the hands, ankles, or feet.
- Anticonvulsants may trigger depression, thoughts of suicide, or other mood disorders in a small percentage of persons who take them for any reason. If you or a family member/caregiver observe any unusual changes in your mood, call your doctor immediately.

Tips for Using Gabapentin Tablet

- To avoid dizziness or sleepiness during the day, take it at the same time each day, ideally before bedtime.
- Antacids should be avoided 2 hours after or before taking Gabapentin Tablet 300 mg because they make it more difficult for your body to digest the medication.
- It has the potential to make you sleepy. Do not do anything else that requires attention until you have a better understanding of how it impacts you.
- If you notice any odd changes in your mood or behaviour, sudden or worsened depression, or thoughts of suicide or conduct, contact your doctor.
- Stopping it suddenly without consulting your doctor may cause seizures to become more frequent.

Gabapentin Tablet 300 mg Supplier

Healthiza is one of the leading Gabapentin Tablet 300 mg supplier and manufacturer, offering the best quality medicine globally through a validated cold chain secure shipment process. We use this process to manufacture, store and deliver medicines at a very specific temperature. Customers can avail of this premium quality pharma product at an affordable price from us.



GET IN TOUCH HEALTHIZA LIFESCIENCE PVT LTD

EMAIL : info@healthizalife.com | sales@healthizalife.com

PHONE : +91- 93289 40307

ADDRESS : A-907, Signature II, Sarkhej 382210 Ahmedabad, India.

WEBSITE : www.healthizalife.com