

Oxybutynin Tablet 2 mg

Product Information :

Pharmaceutical form : Tablet

Category : Prescription Medicine

Pack : 1x10

Brand name : Oxybutynin

Content : Oxybutynin Tablets 2 mg

Therapeutic Use : To treat overactive bladder



Product Description :

What is Oxybutynin Tablet 2 mg?

Oxybutynin tablet 2 mg is used in the treatment of hyperactive bladder in adults and children (bladder muscles contract uncontrollably, causing frequent urine, an urgent need to urinate, and difficulty to control urination). Patients with spina bifida (when the spinal cord does not work correctly before birth) or other nervous system problems that affect the bladder muscles can take oxybutynin as an extended release tablet to regulate bladder muscles. Oxybutynin belongs to the anticholinergic/antimuscarinic drug class. It relaxes the muscles of the bladder.

How to use Oxybutynin

- Oxybutynin is available in tablet, syrup, and extended-release (long-acting) tablet form. The tablets and syrup should be taken every two to four hours. Every day, around the same time(s), take oxybutynin. Carefully follow the advice on your medication label. Do not consume more or less of it, or take it more frequently than your doctor has suggested.
- With lots of water or other fluids, take the extended-release tablets whole. The extended-

release tablet should not be split, chewed, or crushed. If you or your kid has trouble swallowing tablets, tell your doctor.

- Your doctor may begin with a modest dose of Oxybutynin Tablet 2 mg and gradually raise it.
- Oxybutynin may help you manage your symptoms. Even though you feel Fine, keep taking it. Stop using oxybutynin tablet without consulting your doctor.
- Within the first two weeks of treatment, you may see some relief in your symptoms. However, getting the maximum benefit of oxybutynin may take 6–8 weeks.

Precaution - Oxybutynin Tablet 2 mg

If you have allergies with oxybutynin, any other medicines, or any of the substances in oxybutynin tablets, extended-release pills, or syrup, inform your doctor immediately.

- Tell your doctor about any prescribed and non-prescribed medications, vitamins, nutraceuticals, and herbal products you're using or planning to use. Your doctor may need to adjust your prescription dosages or keep a close eye on you for adverse effects.
- If you have or have ever had narrow angle glaucoma (a vision-threatening eye condition), any condition that prevents your bladder from releasing completely, or any condition which makes your stomach to empty slowly or incompletely, tell your doctor.
- If you have pregnancy, planning to become pregnant, or breastfeeding, tell your doctor. Call your doctor when you become pregnant during taking oxybutynin.
- If you have ulcerative colitis, gastroesophageal reflux disease, hiatal hernia, hyperthyroidism, myasthenia gravis, Parkinson's disease, dementia, a irregular heartbeat, high blood pressure, benign prostatic hypertrophy, or heart, liver, or kidney illness, notify your doctor.
- If you're 65 or older, talk to your doctor about the concerns and advantage of taking oxybutynin tablets or syrup.
- Tell your doctor or dentist that you are getting Oxybutynin Tablet 2 mg if you are undergoing surgery, including dental surgery.
- You should be aware that this drug can induce drowsiness or impaired vision. Do not

operate machinery or drive a car until you understand how this medicine affects you.

- It's important to note that oxybutynin can make it more difficult for your body to cooled when it gets too heated. If you get a fever or other indicators of heat stroke after being exposed to heat, such as dizziness, nausea, headache, disorientation, or a fast pulse, call your doctor or seek immediate medical assistance.

How it works

Oxybutynin is a muscarinic antagonist. It stops frequent, urgent, or uncontrolled urinating by relaxing the muscles of the urine bladder.

Side Effects of Oxybutynin

The majorities of side effects are minor and will go away as your body adapts to the medication. If they don't go away or you're worried about them, see your doctor.

- It's possible that you'll experience dry mouth, dizziness, drowsiness, vision problems, dry eyes, nausea, vomiting, stomach discomfort, constipation, diarrhea, headache, odd taste in mouth, dry/flushed skin, and weakness. Notify your doctor immediately if any of these side effects persist or worsen.
- Suck on hard candies or ice chips, chew gum, drink a lot of water, or use a saliva substitute to ease dry mouth. Employ artificial tears or other eye fluids to treat dry eyes.
- Eat dietary fibre, drink plenty of water, and exercise to avoid constipation. You may also require the use of a laxative.
- Remember that your doctor recommended Oxybutynin Tablet 2 mg because he or she believes the advantage to you exceeds the risk of adverse effects. Many persons who use this medicine have no major adverse effects.
- Reduced sexual activity, trouble urinating, rapid heartbeat, indications of kidney infection, changes in mood (confusion), swelling of arms/legs, vision problems (eye pain), seizures, stomach/intestinal blockage should all be reported to your doctor immediately.

Some Tips for Using Oxybutynin Tablet

- Oxybutynin Tablet 2 mg relieves the effects of an overactive bladder by reducing urine urges and regulating urination.

- Caffeine, alcohol, and carbonated beverages can increase your symptoms.
- It may cause dizziness and vision blurring. Don't drive vehicle or do anything else that involves mental focus until you've experienced the effects.
- Constipation, dry lips, and dry eyes are all possible adverse effects. Maintain proper oral hygiene, drink plenty of water, eat sugarless gum, and use caution when wearing contact lenses.
- It may take up to four weeks before you notice any changes in your symptoms. Continue to take it as directed.

Oxybutynin Tablet 2 mg Supplier

Healthiza is one of the leading Oxybutynin Tablet 2 mg supplier and manufacturer, offering the best quality medicine globally through a validated cold chain secure shipment process. We use this process to manufacture, store and deliver medicines at a very specific temperature. Customers can avail of this premium quality pharma product at an affordable price from us.



GET IN TOUCH HEALTHIZA LIFESCIENCE PVT LTD

EMAIL : info@healthizalife.com | sales@healthizalife.com

PHONE : +91- 93289 40307

ADDRESS : A-907, Signature II, Sarkhej 382210 Ahmedabad, India.

WEBSITE : www.healthizalife.com